



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

These last three follow at once from the fact that for the pentagon, quadrilateral, and triangle, we have three, two, and one term in the expression for the area, while the power of  $\pi$  is for the pentagon  $\pi^4$ , for the quadrilateral  $\pi^3$ , for the triangle  $\pi^2$ , in the denominator.

139. Proposed by L. C. WALKER, A. M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

Four points are taken at random on the surface of a given sphere; find the average volume of the tetrahedron formed by the planes passing through the points taken three and three.

Remark by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

No. 139 is the same as No. 130 for which I have sent a solution previously.

140. Proposed by L. C. WALKER, A. M., Professor of Mathematics, Colorado School of Mines, Golden, Col.

Obtain the average area of a triangle formed by a tangent to the four-cusped hypocycloid and the coordinate axes.

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let  $a$ =radius of fixed circle. Then portion of tangent intercepted by coordinate axes= $a$ . Area of triangle= $\frac{1}{2}xy$ , subject to the condition  $x^2 + y^2 = a^2$ .

$$\therefore \text{Average area} = \frac{1}{2} \int_0^a x \sqrt{a^2 - x^2} dx / \int_0^a dx = \frac{1}{6} a^2.$$


141. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, O.

Upon a circular table, radius  $r$ , a variable square plate is thrown at random. What is the probability that the plate will lie wholly on the table?

Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let  $O$  be the center of the given circle, radius  $r$ ;  $ABCD$  the square, center  $E$ ;  $AB=2x$ ,  $OE=z$ ,  $\angle OAE=\theta$ . Then  $AE=x\sqrt{2}$ .

If the center of the square,  $E$ , falls on a circle center  $O$  and radius  $(r-x\sqrt{2})$ , the square will be wholly on the table. If  $E$  falls on a circle, center  $O$  and radius  $z$ , the plate will lie wholly on the table.  $z = \sqrt{r^2 + 2x^2 - 2rx\sqrt{2}\cos\theta}$ . The limits of  $x$  are 0 and  $\frac{1}{2}r\sqrt{2}$ ; of  $\theta$ , 0 and  $\frac{1}{4}\pi - \sin^{-1}(x/r) = \theta'$ . Let  $p$ =chance. Since the whole number of ways  $E$  can fall on the circle is  $\pi r^2$ , we get,



$$p = \frac{\pi \int_0^{\frac{1}{2}r\sqrt{2}} [r - x\sqrt{2}]^2 dx}{\pi r^2 \int_0^{\frac{1}{2}r\sqrt{2}} dx} + \frac{\pi \int_0^{\frac{1}{2}r\sqrt{2}} \int_0^{\theta'} z^2 dx d\theta}{\pi r^2 \int_0^{\frac{1}{2}r\sqrt{2}} \int_0^{\theta'} dx d\theta}$$